

APPENDIX.

SOUTH AUSTRALIA.

REPORT ON THE DETERMINATION OF THE BOUNDARY LINE OF THE COLONIES OF SOUTH AUSTRALIA AND NEW SOUTH WALES, BY CHARLES TODD, F.R.A.S., OBSERVER AND SUPERINTENDENT OF TELEGRAPHS, SOUTH AUSTRALIA.

Observatory and Telegraph Department, Office of Superintendent, Adelaide, 14th December 1868.

SIR,

Having, in compliance with the instructions of the Government, completed the necessary astronomical observations, in conjunction with the Government Astronomer at Sydney, Mr. Smalley, for fixing the position of the common boundary line of South Australia and New South Wales, I have now the honor, herewith, to furnish a joint declaration of the same, signed by Mr. G. R. Smalley on behalf of the Government of New South Wales, and by myself on behalf of the Government of South Australia, together with the following detailed report, and map showing the exact position of the said boundary line, and the relative position, at its northern extremity, of the present boundary line of South Australia and Victoria.

By Imperial legislation the eastern boundary line of South Australia is defined to be the 141st meridian of east longitude.

The existing boundary line between South Australia and Victoria was fixed from observations made by Mr. C. J. Tyers, in 1839.

First. By triangulation with Melbourne.

Second. By chronometric measurement from Sydney.

Third. By lunar observations with sextant near the assumed boundary.

Mr. Tyers made the longitude of the Sandhill on the coast, from which the line starts, to be $141^{\circ} 2' 54''$ east, or about $1\frac{1}{2}$ miles to the west of the mouth of the river Glenelg.

The observations were subsequently checked by Mr. Owen Stanley, F.R.A.S., Commander of H.M.S. *Britomart*, who made the longitude of the same Sandhill $141^{\circ} 2' 21''$ east, by triangulation, and $141^{\circ} 2' 50''$ by chronometer.

The line, taking the Sandhill before referred to as an initial point, was run from the coast ($38^{\circ} 4' 3''$ S. lat. of high water mark) to the Tatiara, or to about 36° S., in 1846-7, and continued to the Murray in 1849-50, a conical pile of stones, eight feet high, being built about $2\frac{1}{2}$ miles south of where the line strikes the river, and another pile twelve feet high, on the most elevated ground, forty-seven chains fifty links south of the river, (vide map appended to this report). This line was proclaimed in the *Government Gazette* of 23rd December 1847.

In the month of March last, by direction of the Government, I visited Sydney to confer with Mr. Smalley, the Government Astronomer of New South Wales, as to the best plan of giving effect to the wishes of the two Governments for defining the boundary line north of the Murray, it having been decided not to accept the line south of the river without first verifying its accuracy, inasmuch as the electric telegraph connecting the observatories of the three colonies afforded a means of determining the 141st degree of east longitude with greater accuracy than was possible when the boundary of South Australia and Victoria was adopted in 1847.

After consultation we agreed—

First. To make, with the co-operation of Mr. Ellery, the Government Astronomer of Victoria, a careful determination of the difference of longitude between the observatories at Sydney and Melbourne by means of the electric telegraph.

Second. To adopt as a basis a certain assumed longitude of the Sydney Observatory from which the boundary line should be measured, viz., the arithmetical mean of the longitude deduced from the present assumed longitude of Melbourne, the difference of longitudes having been ascertained, and that adduced by Mr. E. J. Stone, First Assistant Astronomer at the Royal Observatory, Greenwich, from observations of the moon at Sydney in 1859-60.

Third. That the personal equations between the different observers should be determined, Mr. Todd observing, for that purpose, with Messrs. Smalley and Russell, at Sydney, and with Messrs. Ellery and White, at Melbourne.

Fourth. That Mr. Todd should then proceed to the boundary and erect a transit instrument on the north of the river, near to the line of electric telegraph connecting Adelaide and Sydney, the wires being brought into the temporary observatory, and that its longitude should be determined—

a By voltaic signals exchanged with Sydney on two or more clear nights. The signals to be transits over the meridian, or rather over the several wires of the telescope, of certain stars, previously selected, observed at both places, and recorded on the Sydney chronograph.

b By voltaic signals exchanged in the same way with the Melbourne Observatory on two or more clear nights. The transits being recorded on the Melbourne chronograph.*

Fifth. That the geographical latitude of the transit instrument should then be ascertained by transits of stars over the prime vertical.

Sixth. That the longitude and latitude of the transit instrument having been determined and its distance, east or west, from the 141st meridian mutually agreed to, that distance to be carefully chained, and the boundary line so ascertained to be properly marked and run for a short length for the guidance of the surveyors. A post to be set up on the site of the transit instrument for future reference.

* There being no convenient means of fitting up a chronograph at the boundary, the signals could be sent only one way.

With regard to articles 1 and 2, it may be well to explain that a discrepancy of about three-quarters of a mile was known to exist in the assumed longitudes of the Sydney and Melbourne observatories, and that the position of the 141st meridian would therefore differ by that much according as one or the other were taken as a basis. The agreement arrived at by Mr. Smalley and myself was to divide this difference, but before finally adopting this course we separately recommended and obtained the official concurrence of our respective Governments in the arrangement proposed.

The longitude of the Sydney Observatory, as calculated by Mr. Stone (Royal Astronomical Society's Monthly Notice for June 1867, No. 8, vol. xxvii.), viz., 10h. 4m. 47-32s., is deduced from twenty-four transits of the moon in 1859, and twenty-four in 1860, using only those observations when a transit of the moon was observed at Greenwich not differing more than 40 mins. in right ascension, and where each observation was accompanied by at least one moon culminating star common to both Sydney and Greenwich.

The resulting longitude may be considered free, therefore, from the errors in Burckhardt's lunar tables, which were somewhat large in 1859 and 1860.

Taking each limb separately, the longitude obtained was as follows:—

								h. m. s. Weight.
1859.	From 16 observations of	1st limb	10	4	45-05	38
"	" 8	"	2nd "	...	10	4	48-08	27
1860.	" 13	"	1st "	...	10	4	45-81	47
"	" 11	"	2nd "	...	10	4	50-05	40

Combining the two years we have—

h. m. s.
10 4 45-39 from 29 observations of 1st limb
10 4 49-26 " 19 " 2nd "

And the resulting longitude is 10h. 4m. 47-32s.

A rough comparison by voltaic signals with the observatory at Williamstown, in 1861, made it 10h. 4m. 50-19s., adopting the assumed longitude of Williamstown, 9h. 39m. 38-81s.

The longitude of the observatory at Williamstown, from which that of the new observatory at Melbourne was obtained by a very good triangulation, was deduced from 142 meridian transits of the moon compared with corresponding transits at Greenwich and the Cape of Good Hope, as follows:—

From 89 observations of Moon's 1st limb	h. m. s.
" 62 " 2nd "	9 39 38-86 38-74

Giving weights corresponding to the number of observations, we have as the resulting longitude, 9h. 39m. 38-81s. $\pm 0-19s.$

The difference of longitude between the old observatory at Williamstown and the new one at Melbourne was found by a careful triangulation to be 16-0 secs.; the longitude of the latter is therefore assumed to be 9h. 39m. 54-8s.

Mr. Ellery having kindly promised his cordial co-operation, arrangements were made with Messrs. Cracknell and McGowan for the use of the telegraph wires, and on 3rd April, the night being clear at Sydney, but partially clouded at times at Melbourne, the two observatories were connected in direct circuit, and transits of fifteen stars, previously selected, were observed on both meridians and recorded on the Melbourne chronograph, Messrs. Smalley, Todd, and Russell, observing, in turn, at Sydney, and Mr. E. J. White, the Assistant Astronomer, at Melbourne.

The signals transmitted from Sydney were received by a repeater at Melbourne, the armature of which automatically repeated them to the chronograph. From several trials, Mr. Ellery found that the time lost in repeating was 0-027 secs., which has consequently been subtracted from the time by the Melbourne clock of each Sydney transit to obtain the true sidereal interval.

To eliminate the time occupied in the transmission of the voltaic signals between the two observatories, the Melbourne clock was made to transmit its beats automatically to the Sydney chronograph, where they were compared with the transit clock there.

The transits were all reduced to one observer—Mr. White—as a standard, the personal equations being determined in the following manner:—

From transits of clock stars observed by Messrs. Smalley, Russell, and Todd, on 3rd April, recorded on the Sydney chronograph, the clock at 11h. 12m. (vide Table III. in the Appendix) was 10-979s. slow by Mr. Smalley (S); 10-883s. by Mr. Todd (T); and 10-730s. by Mr. Russell (R). From which it appears that S sees a star on the wire of the telescope, or observes earlier than T by 0-096s., and earlier than R by 0-249s. Hence—

$$\begin{aligned} S-T &= 0-096 \\ S-R &= 0-249 \\ T-R &= 0-153 \end{aligned}$$

Mr. Smalley observed with Mr. White (W) on 13th April 1867, the personal equation being—

$$S-W = 0-302$$

T observed with W in July 1866, and again on 12th and 14th April 1868, the personal equation being—

$$\begin{aligned} T-W &= 0-255, \text{July, 1866} \\ &= 0-234, \text{12th April 1868} \\ &= 0-237, \text{14th April 1868} \end{aligned}$$

For R—W we have—

$$\begin{aligned} T-W &= 0-235 \text{ (adopted)} \\ \text{and } T-R &= 0-153 \\ \therefore R-W &= 0-082 \end{aligned}$$

The personal equations adopted in the following calculations are—

$$\begin{aligned} S-W &= 0-302 \\ T-W &= 0-235 \\ R-W &= 0-082 \end{aligned}$$

The following table shows the difference of longitude between Melbourne and Sydney, deduced from transits of the same stars over the two meridians recorded on the Melbourne chronograph.

DIFFERENCE of Longitude between Melbourne and Sydney.

Star.	Observer at Sydney.	Time by Melbourne Clock at						Sidereal Interval by Melbourne Clock.					
		Sydney Transit.			Sydney Transit + P Equation with W.		Melbourne Transit, (Observer W.)						
		H.	M.	S.	H.	M.	S.	H.	M.	S.	H.	M.	
α Leonis	...	R	9	36	17.454	9	36	17.536	10	1	13.237	24	55.701
γ Leonis	...	S	9	47	33.267	9	47	33.569	10	12	34.275	55.705	
ρ Leonis	...	T	10	0	48.414	10	0	48.619	10	25	44.377	55.728	
β Leonis	...	R	10	17	16.029	10	17	16.111	10	42	11.688	55.777	
B.A.C. 3822	...	S	10	38	20.692	10	38	20.994	11	3	26.456	55.162	
δ Crateris	...	T	10	47	41.763	10	47	41.998	11	12	37.852	55.834	
π Leonis	...	R	11	5	8.631	11	5	8.703	11	30	4.577	55.874	
β Leonis	...	S	11	17	16.247	11	17	16.549	11	42	12.577	55.028	
B.A.C. 4046	...	T	11	27	46.483	11	27	46.718	11	52	42.290	55.572	
B.A.C. 4088	...	R	11	36	29.956	11	36	30.038	12	1	29.741	55.703	
β Corvi	...	T	12	2	25.119	12	2	25.254	12	27	21.100	55.746	
γ Virginis	...	R	12	9	55.575	12	9	55.657	12	34	51.463	55.806	
B.A.C. 4309	...	S	12	18	29.887	12	18	30.189	12	43	25.701	55.512	
B.A.C. 4355	...	T	12	28	17.262	12	28	17.497	12	53	15.319	55.823	
B.A.C. 4383	...	R	12	34	31.922	12	34	32.004	12	59	27.794	55.790	
												15) 11.063	
												24 55.738	
Mean ...												+ .012	
Add for losing rate (0.71s. in 24 hours) of clock												+ .027	
Add for time lost by repeating register													
Difference of longitude, less time occupied by voltaic current ...												24 55.777	

By comparison of clocks—beats of Melbourne clock transmitted to Sydney chronograph—

Time.	Clock slow.	Observer.	1st Comparison.			2nd Comparison.		
			H.	M.	S.	H.	M.	S.
Time by Sydney clock	S	9	48	53.33
Time by Melbourne clock	R	9	24	0.00

Each comparison being derived from the mean of several beats of the two clocks as recorded on the chronograph.

For the error of the Sydney clock by W, at these times, we have—

Time.	Clock slow.	Observer.	Present Equation with W.			Clock slow by W.
			H.	M.	S.	
9 38	...	S	11.05	10.748
9 33	...	T	10.98	10.745
10 1	...	R	10.75	10.688
10 13	...	S	10.99	10.738
10 26	...	T	10.98	10.745
10 42	...	R	10.82	10.688
11 7	...	S	10.95	10.648
11 12	...	T	10.85	10.638
11 30	...	R	10.71	10.788
11 43	...	S	11.09	10.888
12 35	...	R	10.64	10.688
Mean ...	10 49	Sydney clock slow by W	10 688

The daily gaining rate of the clock was 1.50 sec. The Sydney clock was therefore slow at 1st comparison (9h. 49m.) by W 10.694 secs., and 10.685 secs. at 2nd comparison. The Melbourne clock was slow by W at the same times 8.21s. and 8.26s. respectively.

1ST COMPARISON.

1st Time by Sydney clock	Clock slow by W	H. M. S.		
		9	48	53.330
...
...	+	10.694
...

Sidereal time at Sydney ... 10 49 4.024

2nd Time by Melbourne clock	Clock slow by W...	H. M. S.		
		9	24	0.000
...
...	+	8.21
...

Corresponding sidereal time at Melbourne ... 9 24 8.210

Difference of time ... 24 55.814

2ND COMPARISON.

1st Time by Sydney clock	H.	M.	S.
Clock slow by W.	11	18	53.50
			+	10.685		
Sidereal time at Sydney			
2nd Time by Melbourne clock	H.	M.	S.
Clock slow by W.	10	54	0.00
			+	8.25		
Corresponding sidereal time at Melbourne			
Difference of time			
Ditto	(1st Comparison)			
Mean			

Hence—

Difference of longitude (signals transmitted from Sydney to Melbourne)	24	55.777
Ditto ditto (signals transmitted from Melbourne to Sydney)	24	55.874

Difference = twice time occupied by current 0.097

The voltaic current appears, therefore, to have taken 0.048 sec. to pass from one observatory to the other, a distance of about 540 miles, or to have had a velocity of 11,250 miles a second.

Giving twice the weight to the former of the two measurements, we have as the resulting difference of longitude between the two observatories, 24 min. 55.81 sec.

Difference of longitude	H.	M.	S.
Assumed longitude of Melbourne	9	39	54.80
Longitude of Sydney referred to Melbourne	10	4	50.81
Longitude of Sydney by Mr. Stone, Royal Astronomical Society's Notices, No. 8, vol. xxvii.	10	4	47.33
Mean = adopted longitude of Sydney	10	4	48.97

I left Sydney for Adelaide on 7th April, spending a few days at Melbourne to observe with Mr. Ellery and Mr. White, and after a week's stay in Adelaide, I proceeded up the river by the steamer *Prince Alfred* from Blanchetown, accompanied by Mr. A. B. Cooper, Deputy Surveyor-General, taking with me an excellent 4-feet transit instrument, having an aperture of 2½ inches, three chronometers, &c., and arrived at Chowilla on the evening of Friday the 1st of May.

The following day and Monday were occupied in carting instruments and camp to the boundary. A suitable site for the transit instrument was found at a distance of 25 chains 56.68 links to the west of the present boundary between South Australia and Victoria, produced north of the river, and about one-third of a mile from the line of electric telegraph, from which a short line of that length was temporarily run up to bring the wires into the observatory.

The weather being favorable, I was able to get the transit into good adjustment and everything ready for signalling by the following Saturday. And on the nights of 9th and 10th May, the sky being splendidly clear, longitude signals were exchanged with the Sydney Observatory. The circuit (about 900 miles long) was not very good, and many of the boundary transits could not be recorded on the Sydney chronograph, the adjustment being difficult, especially on Sunday night.

The following table shows the resulting difference of longitude corrected for personal equation of observers, Messrs. Smalley and Todd.

TABLE showing the Time by the Transit Clock at Sydney, of Transit of Stars over the meridians of the Sydney Observatory and of Station near the Boundary recorded on the Sydney Chronograph, and the observed Sidereal Interval corrected for personal equation.

Date.	Star observed.	Observer at Sydney.	Observer at Boundary.	Time by Sydney Clock as—			Sidereal Interval by Clock.
				Sydney Transit.	Boundary Transit.		
1863.				H.	M.	S.	
May 9	α Leonis	...	S	T	10	25	48.98
"	B.A.C. 3638	"	10	31	0.41
"	B.A.C. 3926	"	11	26	21.03
"	β Leonis	"	11	42	17.31
"	B.A.C. 4015	"	11	46	13.38
"	B.A.C. 4087	"	11	50	20.38
May 10	β Leonis	"	11	42	15.86
"	B.A.C. 4015	"	11	46	12.21
"	B.A.C. 4087	"	11	50	19.15
"	θ Virginis	"	13	3	3.89
				H.	M.	S.	
				11	6	48.99	41 0.01
				11	11	59.86	40 59.44
				12	7	20.77	40 59.24
				12	23	17.23	40 59.22
				12	27	13.01	40 59.63
				12	31	20.05	40 59.67
				12	35	13.86	41 0.00
				12	37	11.70	40 59.49
				12	31	18.91	40 59.76
				13	44	3.98	41 0.09
							10) 7.75
				Mean	
	Clock's loss in 41 min.	40 59.75
	Personal equation, T - S	+ .009
					- .006
	Difference of longitude	40 59.718

On 13th and 14th May, signals were successfully exchanged between the boundary and Melbourne, Messrs. Cracknell and McGowan having kindly arranged to give me direct circuit with the Melbourne Observatory, via Deniliquin and Echoes, all intermediate telegraph offices being cut out. The nights were brilliantly clear at both places, and the circuit splendid.

Zenith stars were selected, nine stars being observed on 13th May and twelve on 14th May. On the second night the transit at the boundary was reversed in the middle of the series, six stars being taken with lamp east, and six with lamp west. B.A.C. 655 S.P. and B.A.C. 4790 were observed for azimuth error at the boundary on both nights, the transits being recorded on the Melbourne chronograph.

The following two tables show the resulting difference of longitude between the Melbourne Observatory and Boundary Transit, corrected for personal equation of observers ($T - W = 0.235$ seconds).

1868.—13TH MAY.

Name of Star.	Time by Melbourne Clock of Transit at—						Difference.	
	Melbourne. (W)			Boundary. (T)				
	H.	M.	S.	H.	M.	S.		
B.A.C. 4035	12	2	41.202	12	18	45.150	16 3.928	
α Centauri	12	16	6.755	12	32	10.495	16 3.740	
B.A.C. 4223	12	30	9.246	12	46	13.538	16 3.532	
B.A.C. 4329	12	42	59.541	12	59	2.813	16 3.512	
B.A.C. 4325	12	52	46.275	13	8	59.580	16 3.605	
B.A.C. 4417	13	4	9.745	13	20	13.253	16 3.538	
β Centauri	13	12	38.892	13	28	42.363	16 3.471	
B.A.C. 4507	13	22	51.825	13	38	55.453	16 3.628	
ι Centauri	13	37	39.411	13	53	42.736	16 3.325	
Personal equation, $T - W$	Mean	16 3.508	
Transmission time	+ .235	
Correction for repeater	- .026	
Difference of longitude from nine stars on 13th May	- .927	
							16 3.780	

14TH MAY.

Name of Star.	Time by Melbourne Clock of Transit at—						Difference.	
	Melbourne. (W)			Boundary. (T)				
	H.	M.	S.	H.	M.	S.		
B.A.C. 4046	11	52	42.670	12	8	53.363	16 3.693	
B.A.C. 4095	12	3	15.056	12	19	18.652	16 3.596	
α Centauri	12	16	40.714	12	32	44.202	16 3.488	
B.A.C. 4202	12	21	23.558	12	37	26.859	16 3.501	
B.A.C. 4253	12	30	43.787	12	46	47.379	16 3.592	
B.A.C. 4309	12	43	33.270	12	52	36.855	16 3.585	
B.A.C. 4355	12	55	20.305	13	9	24.377	16 3.442	
B.A.C. 4417	13	4	43.653	13	20	47.253	16 3.810	
ι Centauri	13	13	12.794	13	29	16.173	16 3.449	
B.A.C. 4507	13	23	25.709	13	39	29.386	16 3.677	
ι Centauri	13	38	13.185	13	54	16.999	16 3.814	
κ Centauri	13	44	14.662	14	0	18.126	16 3.664	
Personal equation, $T - W$	Mean	16 3.576	
Transmission time	+ .235	
Correction for repeater	- .026	
Difference of longitude from twelve stars	- .027	
							16 3.758	

Difference on 13th May = 16 3.780, Weight 9.

Difference on 14th May = 16 3.758, Weight 12.

Hence the difference of longitude between the observatory at Melbourne and transit near boundary, was 16 min. 3.767 sec.; and for resulting longitude of the transit instrument we have—

1ST.

Adopted longitude of Sydney 10 4 48.97
Difference of longitude between transit and Sydney 40 59.72

Longitude of transit by signals with Sydney 9 23 49.25

2ND.

Difference of longitude between transit and Melbourne 16 3.77

Difference of longitude between Sydney and Melbourne 24 55.81

Sum = difference of longitude between transit and Sydney 40 59.58

Adopted longitude of Sydney 10 4 48.97

Longitude of transit by signals with Melbourne 9 23 49.39

The mean of these is 9 hours 23 min. 49.32 secs. ($140^{\circ} 57' 19.8''$ E.), or, giving to each determination a weight proportionate to the number of stars observed, 9 hours 23 min. 49.34 secs.

The longitude of the transit instrument actually adopted, from calculations made on the spot, and from which the boundary line was measured, was 9 hours 23 min. 49.312 secs. ($140^{\circ} 57' 19.7''$), which is practically the same.

To obtain the latitude the piers were shifted on 20th May and transits of stars over the prime vertical were observed on that and the following night; care being taken to have the centre of the transit instrument exactly over the same spot as before.

Only one star, No. 4603 in the British Association's catalogue, was observed on both sides of the meridian on 20th May; but on the following night B.A.C. 4437, κ^2 Centauri, B.A.C. 4603, and B.A.C. 4916 were taken at both transits.

The adopted latitude from these, deduced at the time of transits east and west, over the middle wire only, was $33^{\circ} 55' 8''$, which may be a few seconds in error, the apparent places of the stars being calculated from the British Association's catalogue, and requiring verification.

The adopted position of the transit instrument being $140^{\circ} 57' 19.7''$ east longitude, and $33^{\circ} 55' 8''$ south latitude, its distance due west from the 141st meridian was therefore 2 miles 44 chains 68½ links.

Mr. Smalley, on behalf of the New South Wales Government, having telegraphed to me his acceptance of this result, the distance just mentioned was carefully chained on a line due east from the centre of the transit instrument by Mr. Cooper, and subsequently, as a check, by myself.

The 141st meridian was then carefully run south to where it intersects the upper and lower tracks to Wentworth, or for nearly two miles.

A stout post, painted white, was planted on the exact site of the transit instrument, with the following painted, in large black letters and figures, viz. —

On south face of pole :

Longitude, $140^{\circ} 57' 19.7''$ East. Latitude, $33^{\circ} 55' 8''$ South.

On east face :

↑ C. Todd, Observer.

On north face :

Province Boundary, 2 miles 44 chains 68 links East.

On west face :

↑


On the boundary line, where it crosses the upper and lower tracks to Wentworth, a stout white painted post was planted on each side of the two tracks, lettered on the side facing the track —

"PROVINCE BOUNDARY;"

on the east side the letters "N. S. W.;" and on the west side, "S. A."

Mr. Smalley met me, by appointment, at Wentworth last month, and proceeded with me to the boundary, where he formally accepted it, as marked on the ground, on behalf of the Government of New South Wales.

The position of the boundary thus determined from the foregoing observations, and mutually agreed to by Mr. Smalley and myself on behalf of our respective Governments, would be defined as a meridian line starting from the river Murray, 2 miles 19 chains due east of the pile of stones on the south bank of the river (mentioned in a former part of this report), marking the north end of the line at present adopted as the common boundary of South Australia and Victoria; as, however, there were no good natural landmarks we thought it better to permanently indicate the exact position of the new boundary line by a brick pyramid, 13 feet 6 inches high and 5 feet 6 inches square at the base, which we erected on the slope of the scarp (vide point A in accompanying map), forming the limit of the Murray floods, and immediately to the north of a circular salt lagoon (dry at the time), having the words "PROVINCE BOUNDARY" on the north and south faces; "N. S. W." } on the east face, and "S. A." } on the west face.



The above mark is erected a short distance up the slope of the scarp, about 77 yards from the nearest point of the line of electric telegraph, on the north side of the line. The bend of the river Murray, immediately to the east of Slaney's Island, being nearly $3\frac{1}{2}$ miles distant on an astronomical bearing of about 53° west of south; the Salt Creek public-house being 2 miles 71 chains distant, 30° east of south; and Mount Hancock bearing about $16\frac{1}{2}^{\circ}$ east of south.

During my stay at the boundary, Mr. Cooper assisted me in making a series of magnetic observations with a view of determining the declination of the compass, and dip.

The resulting declination at different hours of the day is shown in the following table:—

Hours of Observation.	Number of Observations.	Declination.
7 a.m. to 9 a.m.	...	6 31 18 east
9 " to 11 "	7	6 34 46 "
11 " to 12 noon	3	6 34 53 "
12 noon to 1 p.m.	3	6 37 29 "
1 p.m. to 3 "	7	6 39 32 "
3 " to 5 "	4	6 36 3 "
5 " to 8 "	7	6 33 15 "

Adopted mean variation, $6^{\circ} 35' 15''$ east. The magnetic dip, from eleven observations between the same dates, was found to be $64^{\circ} 5' 7''$.

In conclusion, I have to gratefully acknowledge the cordial manner in which Mr. Smalley, acting on behalf of the New South Wales Government, co-operated with me throughout. From both Mr. Smalley and his assistant, Mr. Russell, as well as, indeed, from Mr. Cracknell, and many other scientific friends, I received during my stay in Sydney the kindest attention. Nor should I omit to acknowledge the valuable assistance of Mr. Ellery and the Assistant Astronomer, Mr. White, whose co-operation these pages will have shown was essential to the successful completion of the work entrusted to me.

Mr. Cracknell and Mr. McGowan gave every facility for the free use of the lines of electric telegraph under their respective control. I have also to thank Mr. Cooper for his ever ready and zealous aid in the somewhat arduous labors at the boundary.

I have the honor, &c.,

CHARLES TODD,

Observer and Superintendent of Telegraphs.

The Honorable the Treasurer.

APPENDIX.

OBSEVATIONS FOR DETERMINING THE DIFFERENCE OF LONGITUDE BETWEEN THE OBSERVATORIES AT SYDNEY AND MELBOURNE, 3RD APRIL 1868.

TABLE I.—Transits observed at Sydney, and recorded on the Chronograph at the Melbourne Observatory.

Name of Star.	Concluded Transit over Seven Wires.	c. sec. δ	n. tan. δ	m.	Time by Melbourne Clock of true Transit at Sydney.	Observer.
α Leonis	...	9 13 16:087	+ 0:893	- 0:034	+ 0:976	9 13 17:922 S
π Leonis	...	9 28 9:540	+ 0:833	- 0:011	9 28 11:128 T	
α Leonis	...	9 36 13:662	+ 0:833	- 0:017	9 36 17:444 R	
γ Leonis	...	9 47 36:450	+ 0:869	- 0:028	9 47 38:267 S	
ρ Leonis	...	10 0 46:626	+ 0:825	- 0:013	10 0 48:414 T	
ι Leonis	...	10 17 14:239	+ 0:829	- 0:015	10 17 16:029 R	
B.A.C. 3783	...	10 31 54:526	+ 0:931	+ 0:045	10 31 56:498 T	
B.A.C. 3822	...	10 38 23:714	+ 0:936	+ 0:046	10 38 30:692 S	
δ Leonis	...	10 42 0:170	+ 0:872	- 0:029	10 42 1:989 S	
δ Hydra et Crateris	...	10 47 39:931	+ 0:838	+ 0:018	10 47 41:763 T	
ν Leonis	...	11 5 6:831	+ 0:813	+ 0:001	11 5 8:621 R	
β Leonis	...	11 17 14:449	+ 0:843	- 0:021	11 17 16:247 S	
B.A.C. 4046	...	11 27 44:467	+ 0:988	+ 0:052	11 27 46:483 T	
B.A.C. 4088	...	11 36 27:949	+ 0:980	+ 0:051	11 36 29:956 R	
β Chamaeleontis	...	11 45 38:959	+ 4:106	+ 0:371	11 45 44:412 S	
β Hydri, S.P.	...	11 53 37:294	- 3:910	- 0:353	11 53 34:907 S	
β Corvi	...	12 2 23:230	+ 0:882	+ 0:031	12 2 23:119 T	
γ Virgilia (1st Star)	...	12 9 53:776	+ 0:813	+ 0:010	12 9 55:575 R	
B.A.C. 4309	...	12 18 27:889	+ 0:973	+ 0:049	12 18 29:887 S	
B.A.C. 4355	...	12 28 15:270	+ 0:968	+ 0:048	12 28 17:262 T	
B.A.C. 4383	...	12 34 29:898	+ 0:995	+ 0:063	12 34 31:922 R	

$$\text{Correction to meridian} = m + c \text{ sec. } \delta + n \tan. \delta$$

The collimation error (c) = + 0:813s. Level error (δ) = + 0:853s.

The value of n was obtained from transits of β Chamaeleontis and β Hydri S.P.

$$(t^0 - t) - (a^0 - a)$$

$$n = \frac{t^0 - t}{a^0 - a} = 0:075s.$$

$\tan. \delta - \tan. \beta$,

Where—

t = observed time of transit of β Chamaeleontis; corrected for collimation error.

t^0 = observed time of transit of β Hydri S.P.; corrected for collimation error.

a = apparent R.A. of β Chamaeleontis 12h. 10m. 48:70s. *

a^0 = apparent R.A. of β Hydri ob. 18m. 35:29s. + 12h.†

δ = apparent declination of β Chamaeleontis.

β = apparent declination of β Hydri S.P.

m = b sec. ϕ = n tan. ϕ , where ϕ = latitude $35^{\circ} 51' 41''$.

* Apparent R.A. by nautical almanac has been increased by 0:38 sec. (the Melbourne correction).

† Apparent R.A. by N.A. diminished by 0:10 sec. (the Melbourne correction).

DIFFERENCE OF LONGITUDE BETWEEN SYDNEY AND MELBOURNE.

TABLE II.—Transits observed at Melbourne, and recorded on the Chronograph at the Melbourne Observatory, 3rd April 1868.

Name of Star.	Concluded Transit over Seven Wines.			c. sec. δ	α tan. δ	π	Time by Melbourne Clock of true Transit at Melbourne.			Observer.
α Leonis	11. 10. 1. 13.281	- 0.009	+ 0.049	- 0.084	10. 1. 13.237	10. 1. 13.237	W
γ Leonis	10. 12. 34.287	- 0.010	+ 0.082		10. 12. 34.275		
ρ Leonis	10. 25. 44.431	- 0.009	+ 0.039		10. 25. 44.377		
ι Leonis	10. 42. 11.937	- 0.009	+ 0.044		10. 42. 11.888		
B.A.C. 3829	11. 5. 26.686	- 0.011	- 0.135		11. 5. 26.456		
τ Octantis S.P.	11. 5. 42.700	+ 0.087	+ 7.009		11. 5. 49.912		
δ Hydræ et Crateris	11. 12. 37.980	- 0.009	- 0.055		11. 12. 37.832		
ν Leonis	11. 30. 4.670	- 0.009	0.000		11. 30. 4.577		
Lacaille 4865	11. 34. 36.180	- 0.009	- 2.398		11. 34. 33.599		
β Leonis	11. 42. 12.610	- 0.009	+ 0.060		11. 42. 12.577		
B.A.C. 4046	11. 52. 42.537	- 0.011	- 0.132		11. 52. 42.290		
B.A.C. 4088	12. 1. 25.984	- 0.011	- 0.148		12. 1. 25.741		
β Hydræ. S.P.	12. 18. 28.960	+ 0.043	+ 1.035		12. 18. 29.954		
β Corvi	12. 27. 21.286	- 0.010	- 0.092		12. 27. 21.100		
γ Virginis (1st Star)	12. 34. 51.539	- 0.009	- 0.063		12. 34. 51.463		
B.A.C. 4309	12. 43. 25.940	- 0.011	- 0.144		12. 43. 25.701		
B.A.C. 4355	12. 53. 13.556	- 0.011	- 0.142		12. 53. 13.319		
B.A.C. 4383	12. 59. 28.944	- 0.011	- 0.155		12. 59. 27.794	u	

Correction to meridian = $\alpha + c \sec. \delta \pm \tan. \delta$

Collimation error (c) = - 0.009s.; level error = - 0.20s.

Apparent R.A. of τ Octantis S.P. = 23h. 3m. 58.11s. + 12h.

Apparent R.A. of Lacaille 4865 = 11h. 34m. 41.81s.

 $\alpha = - 0.220s.$

Latitude of the Melbourne Observatory, 37° 49' 53".

PERSONAL EQUATION.

TABLE III.—Sydney Transits recorded on the Sydney Chronograph for determining the Personal Equation between Messrs. Smalley, Todd, and Russell.

Name of Star.	Concluded Transit over Seven Wines.			c sec. δ	α tan. δ	π	True TIME.	Apparent R.A. by S.A. + Melbourne corrections.			Clock slow.	Observer.
α Leonis	9. 38. 9.35	+ 0.093	- 0.034	+ 0.976	9. 38. 11.18	9. 38. 22.23	11.05	S	
π Leonis	9. 53. 2.48	+ 0.023	- 0.011		9. 53. 4.27	9. 53. 15.25	10.98	T	
α Leonis	10. 1. 8.93	+ 0.033	- 0.017		10. 1. 10.72	10. 1. 21.47	10.75	R	
γ Leonis	10. 12. 29.77	+ 0.069	- 0.028		10. 12. 31.69	10. 12. 42.55	10.99		
ρ Leonis	10. 25. 39.93	+ 0.035	- 0.013		10. 25. 41.72	10. 25. 52.70	10.98	S	
ι Leonis	10. 45. 7.61	+ 0.029	- 0.015		10. 42. 9.40	10. 42. 20.22	10.82	R	
δ Leonis	11. 6. 53.57	+ 0.072	- 0.029		11. 6. 55.39	11. 7. 6.84	10.95	S	
δ Hydræ et Crateris	11. 13. 33.40	+ 0.038	+ 0.018		11. 12. 39.23	11. 12. 46.08	10.85	T	
ν Leonis	11. 30. 0.30	+ 0.013	+ 0.001		11. 30. 2.09	11. 30. 12.80	10.71	R	
β Leonis	11. 42. 7.91	+ 0.043	- 0.021		11. 42. 9.71	11. 42. 20.80	11.09	S	
γ Virginis	12. 34. 47.35	+ 0.013	+ 0.010		12. 34. 49.15	12. 34. 59.79	10.64	R	

Clock Slow.

By T.

By R.

h. m.	s.	h. m.	s.	h. m.	s.	
At 9. 38	11.05		At 9. 33	10.98
10. 13	10.99		10. 26	10.98
11. 7	10.95		11. 13	10.85
11. 42	11.09				

h. m.	s.	h. m.	s.	h. m.	s.	
Mean	10. 40	11.020	10. 39	10.987
Clock's gainning rate in 24 hours = + 1.85s.	0.33	=	- 0.041	0.42	=	- 0.054
11. 12	10.979		11. 12	10.883
11. 12	10.730		11. 12	10.730

Hence S - T = 0.096s.

S - R = 0.249s.

T - R = 0.153s.

DIFFERENCE OF LONGITUDE BETWEEN THE BOUNDARY TRANSIT AND THE SYDNEY OBSERVATORY.

TABLE IV.—Sydney Transits recorded on the Sydney Chronograph, 9th May 1868.

Name of Star.	Concluded Transit over Seven Wires.	Correction for Collimation Error.	Correction for Level Error.	Correction for Azimuth Error.	True Transit.	Observer.
ρ Leonis ..	10 25 47.75	+ 0.85	+ 0.40	- 0.02	10 25 48.98	S
B.A.C. 3638 ..	10 30 58.77	+ 0.94	+ 0.70	0.00	10 31 04.1	"
B.A.C. 3926 ..	11 26 19.33	+ 0.97	+ 0.73	0.00	11 26 21.03	"
β Leonis ..	11 42 16.03	+ 0.87	+ 0.43	- 0.02	11 42 17.31	"
B.A.C. 4015 ..	11 46 11.63	+ 1.00	+ 0.75	0.00	11 46 13.38	"
B.A.C. 4037 ..	11 50 18.64	+ 0.99	+ 0.75	0.00	11 50 20.38	"

TABLE V.—Boundary Transits recorded on the Sydney Chronograph, 9th May 1868.

Name of Star.	Concluded Transit over Seven Wires.	Collimation Error, - 0.002s.	Level Error, + 1.47s.	Azimuth Error, 0.93.	True Transit.	Observer.
ρ Leonis ..	11 6 47.93	0.00	+ 1.06	0.00	11 6 48.99	T
B.A.C. 3638* ..	11 11 58.22	0.00	+ 1.63	0.00	11 11 59.85	"
B.A.C. 3926 ..	12 7 19.07	0.00	+ 1.70	0.00	12 7 20.77	"
β Leonis ..	12 23 16.23	0.00	+ 1.00	0.00	12 23 17.23	"
B.A.C. 4015 ..	12 27 11.26	0.00	+ 1.75	0.00	12 27 13.01	"
B.A.C. 4037 ..	12 31 18.31	0.00	+ 1.74	0.00	12 31 20.05	"

* First four wires (A, B, C, D) not recorded on the Sydney chronograph; correction to mean of wires + 34.31 secs. applied, calculated from the following equatorial intervals, viz.—A = + 40.189 secs.; B = + 30.685 secs.; C = + 15.431 secs.; D = + 9.167 secs.; E = - 13.268 secs.; F = - 30.939 secs.; G = - 66.337 secs.

TABLE VI.—Sydney Transits recorded on the Sydney Chronograph, 10th May 1868.

Name of Star.	Concluded Transit over Seven Wires.	Collimation Error.	Level Error.	Azimuth Error.	True Transit.	Observer.
β Leonis ..	11 42 14.77	+ 0.75	+ 0.50	- 0.16	11 42 15.86	S
B.A.C. 4015 ..	11 46 10.45	+ 0.87	+ 0.89	0.00	11 46 12.21	"
B.A.C. 4037 ..	11 50 17.43	+ 0.85	+ 0.87	0.00	11 50 19.15	"
θ Virginis* ..	12 3 2.62	+ 0.72	+ 0.65	- 0.10	12 3 3.89	"

* The third wire lost, + 1.25s. applied to mean of wires.

TABLE VII.—Boundary Transits recorded on the Sydney Chronograph, 10th May 1868.

Name of Star.	Concluded Transit over Nine Wires.	Collimation Error, + 0.025s.	Level Error, - 0.12s.	Azimuth Error, + 0.448s.	True Transit.	Observer.
β Leonis ..	12 23 16.20	+ 0.09	- 0.08	- 0.35	12 23 15.86	T
B.A.C. 4015 ..	12 27 11.76	+ 0.09	- 0.14	- 0.01	12 27 11.70	"
B.A.C. 4037 ..	12 31 18.97	+ 0.09	- 0.14	- 0.01	12 31 18.91	"
θ Virginis ..	13 44 4.22	+ 0.08	- 0.10	- 0.22	13 44 3.98	"

The transits at the boundary were taken over nine wires, respectively designated A, B, C, I, D, II, E, F, G. The whole of the above transits on 10th May were recorded on the chronograph at Sydney. In all, 38 signals from the boundary were recorded at Sydney on 9th May and 36 on 10th May. Several other stars out of the list selected were observed, but, owing to difficult circuit, only those given in the foregoing tables could be recorded.

For Azimuth Error of Transit at Boundary on 9th and 10th May.

9TH MAY.

β Chamaleontis.
 $\delta 78^{\circ} 35'$
 $ZD 45^{\circ} 27'$
H. M. S.

Wire A	9	41.0
" B	10	58.0
" C	12	16.0
" D	13	33.0
" E	14	51.0
" F	12	16.9.2
" G	17	28.22

β Hydry S.P.
 $\delta 77^{\circ} 39' 35''$
 $ZD 68^{\circ} 52'$
H. M. S.

Wire A	17	51.0
" B	19	3.2
" C	20	16.5
" D	21	29.2
" E	22	43.0
" F	12	23.58.0
" G	12	23.58.0

6) 17 28.22

6) 123 20.9

12 12 34.70

12 20 53.48

Correction 38.89

— 37.00

Mean 12 12 15.81

12 20 16.48

Collin. — 0.002 \times + 5.05 = — 0.01

— 0.002 \times + 4.809 = + 0.01

Level + 1.47 \times + 3.59 = + 5.23

+ 1.47 \times — 1.79 = — 2.63

Azimuth 0.0 \times — 3.55 = 0.00

0.0 \times + 4.46 = 0.00

12 12 21.08

12 20 13.86

$$\text{Azimuth Error} = \frac{(\Lambda' - \Lambda) - (T' - T)}{h' - h} = \frac{0.01}{8.61} = 0.00$$

Where Λ' = apparent right ascension of β Hydry SP + 12 δ

Λ = " β Chamaleontis

T' = observed time of transit of β Hydry SP

T = " β Chamaleontis

h' and h = Azimuth factors (see δ sin ZD).

H. M. S.
 $\Lambda' = 12 18 40.15$
— 10
 $\Lambda = 12 10 46.88$
+ 38

H. M. S.
 $T' = 12 20 13.86$
 $T = 12 12 21.08$

$h' = + 4.46$
 $h = - 3.55$

$\Lambda' - \Lambda = 7 52.79$

$T' - T = 7 52.78$

$h' - h = + 8.01$

10TH MAY.

β Chamaleontis.
H. M. S.
A 8 27.0
B 9 46.0
C 11 3.0
D 12 19.3
E 13 37.0
F 14 56.3
G 12 16 14.0

β Hydry SP.
H. M. S.
— 17 48.0
19 3.0
20 17.0
21 30.0
22 43.3
12 23 58.0

7) 16 22.6

6) 5 19.3

13 12 20.37

12 20 53.22

12 12 21.53

— 37.00

12 12 20.37

12 20 16.22

Collin. — 0.002 \times + 5.05 = — 0.10

— 0.002 \times + 4.809 = + 0.01

Level — 0.12 \times + 3.59 = — 0.43

— 0.12 \times — 1.79 = + 0.21

Azimuth — 0.45 \times — 3.55 = + 1.60

— 0.45 \times — 4.46 = — 2.01

12 12 21.53

12 20 14.43

$$\text{Azimuth Error} = \frac{-3.59}{+8.01} = -0.448.$$

DIFFERENCE OF LONGITUDE BETWEEN BOUNDARY TRANSIT AND MELBOURNE.

TABLE VIII.—Transits observed at Melbourne, and recorded on the Chronograph at Melbourne, 13th May 1868.

Name of Star.	Concluded Transit over Seven Wires.	α sec δ	α tan δ	α	True Transit.	Observed
ν Octantis S.P.	...	10 4 40.487	-0.272	+3.699	-0.130	10 4 43.784
Lacaille 4342	...	10 11 50.576	+0.256	-3.338	10 11 47.344	+
B.A.C. 4095	...	12 3 41.290	+0.019	+0.023	12 2 41.202	+
α Centauri	...	12 16 6.847	+0.020	+0.018	12 16 6.755	+
B.A.C. 4253	...	12 30 9.997	+0.018	+0.061	12 30 9.946	+
B.A.C. 4309	...	12 42 59.426	+0.019	+0.026	12 42 59.341	+
B.A.C. 4355	...	12 52 47.057	+0.019	+0.029	12 52 46.975	+
B.A.C. 4417	...	13 4 9.851	+0.020	+0.004	13 4 9.745	+
α Centauri	...	13 12 38.991	+0.020	+0.011	13 12 38.892	+
B.A.C. 4507	...	13 22 51.939	+0.021	+0.005	13 22 51.825	+
δ Centauri	...	13 37 39.491	+0.019	+0.031	13 37 39.411	+

Collimation error (ϵ) = + 0.0168; level error = - 0.1038.

Apparent R.A. of ν Octantis S.P. = 22h. 5m. 18.70s. + 12h.

Apparent R.A. of Lacaille 4342 = 10h. 12m. 22.25s.

α = - 0.218s.

DIFFERENCE OF LONGITUDE BETWEEN BOUNDARY TRANSIT AND MELBOURNE.

TABLE IX.—Transits observed at the Boundary, and recorded on the Chronograph at Melbourne,
13th May 1868.

Name of Star.	Concluded Transit over Five Wires.			c. sec. δ	n. tan. δ	m.	True Transit.	Observer
B.A.C. 4095	H. M. S.	S.	S.	S.	H. M. S.	T
...	12 18 44.716	+ 0.258	- 0.001	+ 0.157	12 18 45.130	
ζ Centauri	12 32 10.094	+ 0.260	- 0.016		12 32 10.495	
B.A.C. 4253	12 46 12.998	+ 0.289	+ 0.144		12 46 13.538	
B.A.C. 4309	12 59 2.466	+ 0.256	+ 0.014		12 59 2.893	
B.A.C. 4355	13 8 50.144	+ 0.255	+ 0.024		13 8 50.580	
B.A.C. 4417	13 20 12.926	+ 0.268	- 0.048		13 20 13.283	
ι Centauri	13 28 41.986	+ 0.264	- 0.044		13 28 42.363	
B.A.C. 4507	13 38 55.140	+ 0.261	- 0.105		13 38 55.455	
λ Centauri	13 53 42.294	+ 0.253	+ 0.032		13 53 42.736	
B.A.C. 655 S.P.	14 15 4.457	- 1.564	+ 5.945		14 15 8.975	
B.A.C. 4790	14 43 2.850	+ 5.115	- 19.609		14 42 48.493	

Transit over wires C.I.D.I.E.

Collimation error + 0.214s.; level error + 0.130s.

Apparent R.A. of B.A.C. 655 S.P. = 1h. 59m. 41.10s. + 12h. S.P.D. = 7° 51' 44".

Apparent R.A. of B.A.C. 4790 = 14h. 27m. 20.61s. S.P.D. = 2° 23' 52".

s = - 0.821s.

DIFFERENCE OF LONGITUDE BETWEEN THE BOUNDARY TRANSIT AND MELBOURNE.

TABLE X.—Transits observed at Melbourne, and recorded on the Melbourne Chronograph,
14th May 1868.

Name of Star.	Concluded Transit over Seven Wires.			c. sec. δ	n. tan. δ	m.	True Transit.	Observer
η Octantis	H. M. S.	S.	S.	S.	H. M. S.	W
τ Octantis S.P.	11 0 19.827	+ 0.150	- 2.294	- 0.129	11 0 17.554	
B.A.C. 4046	11 6 11.240	- 0.500	+ 7.833		11 6 18.425	
B.A.C. 4095	11 52 49.759	+ 0.019	+ 0.021		11 52 49.670	
ζ Centauri	12 3 15.141	+ 0.019	+ 0.023		12 3 15.036	
B.A.C. 4302	12 16 40.803	+ 0.019	+ 0.021		12 16 40.714	
B.A.C. 4253	12 21 23.470	+ 0.020	- 0.003		12 21 23.358	
B.A.C. 4309	12 30 43.829	+ 0.018	+ 0.019		12 30 43.787	
B.A.C. 4355	12 43 33.350	+ 0.019	+ 0.030		12 43 33.270	
B.A.C. 4417	12 53 21.013	+ 0.019	+ 0.032		12 53 20.935	
ι Centauri	13 4 43.757	+ 0.020	+ 0.003		13 4 43.655	
B.A.C. 4507	13 13 12.821	+ 0.020	+ 0.012		13 13 12.724	
λ Centauri	13 23 23.823	+ 0.021	- 0.006		13 23 25.709	
λ ¹ Centauri	13 38 15.260	+ 0.019	+ 0.035		13 38 15.185	
			13 44 14.737	+ 0.019	+ 0.035		13 44 14.662	

Collimation error (c) = + 0.016s.; level error = - 0.102s.

Apparent R.A. of η Octantis 11h. 0m. 17.76s.

Apparent R.A. of τ Octantis S.P. 23h. 0m. 18.64s. + 12h.

s = - 0.246s.

DIFFERENCE OF LONGITUDE BETWEEN BOUNDARY TRANSIT AND MELBOURNE.

TABLE XI.—Transit observed at the Boundary, and recorded on the Melbourne Chronograph, 14th May 1868.

Name of Star.	Concluded Transit over Five Wires.	c. sec. δ	n. sec. δ	m.	True Transit.	Observer.
B.A.C. 4046	12 8 53:432	+ 0:159	- 0:015	- 0:193	12 8 53:383	T
B.A.C. 4095	12 19 18:708	+ 0:158	- 0:001	12 19 18:673	"	
α Centauri	12 32 44:274	+ 0:159	- 0:018	12 32 44:222	"	
B.A.C. 4202	12 37 27:010	+ 0:167	- 0:015	12 37 26:679	"	
B.A.C. 4253	12 46 47:290	+ 0:146	+ 0:156	12 46 47:399	"	
B.A.C. 4309	12 59 36:895	+ 0:157	+ 0:015	12 59 36:875	"	
*B.A.C. 4355	13 9 24:788	- 0:156	+ 0:026	- 0:261	13 9 24:397	"
B.A.C. 4417	13 20 47:782	- 0:164	- 0:074	13 20 47:283	"	
β Centauri	13 29 16:664	- 0:162	- 0:048	13 29 16:193	"	
B.A.C. 4507	13 39 29:950	- 0:168	- 0:115	13 39 29:406	"	
i Centauri	13 54 17:400	- 0:155	+ 0:035	13 54 17:019	"	
λ ¹ Centauri	14 0 18:326	- 0:155	+ 0:036	14 0 18:146	"	
B.A.C. 655 S.P.	14 15 36:156	+ 0:958	+ 6:446	14 15 43:319	"	
B.A.C. 4790	14 43 47:486	- 3:131	- 21:238	14 43 22:706	"	

* Transit reversed after B.A.C. 4355. Order of wires before reversion, C.I.D.I.L.L.; lamp east. After reversion, E.I.D.I.L.L.; lamp west. Level error, lamp east, - 0:008; lamp west, - 0:217. The values of c. and n. were deduced from the following equations.

$$\begin{aligned}
 3:569 + 1:215c. + 0:017n. &= 3:592 - 1:190c. - 0:029n. \\
 3:459 + 1:206c. + 0:001n. &= 3:868 - 1:254c. + 0:083n. \\
 3:367 + 1:216c. + 0:020n. &= 3:679 - 1:236c. + 0:054n. \\
 3:459 + 1:274c. + 0:118n. &= 3:980 - 1:282c. + 0:129n. \\
 3:310 + 1:117c. - 0:175n. &= 3:954 - 1:184c. - 0:039n. \\
 3:433 + 1:196c. - 0:017n. &= 3:603 - 1:184c. - 0:040n.
 \end{aligned}$$

$$\begin{aligned}
 \text{Sum. } 29:597 + 7:224c. - 0:036n. &= 22:676 - 7:330c. + 0:158n. \\
 14:554c. &= 2:079 + 0:194n. \\
 c. &= + 0:143 + 0:013n.
 \end{aligned}$$

From B.A.C. 655 S.P. and B.A.C. 4790, we have
 $55:006 + 7:310c. - 7:241n. = 86:905 - 23:902c. + 23:584n.$
 $31:212c. = 31:125c. + 31:900.$
 $c. = 0:988n. + 1:022.$
 $0:988n. = - 0:879.$
 $n. = - 0:883.$
 $c. = + 0:131.$

CORRECTIONS APPLIED TO APPARENT RIGHT ASCENSIONS OF NAUTICAL ALMANAC STARS FROM MELBOURNE OBSERVATIONS, 1860 TO 1867.

α Leonis	0:00
π Leonis	+ 0:01
α Leonis	+ 0:02
γ Leonis	- 0:02
β Leonis	- 0:04
λ Leonis	+ 0:02
δ Leonis	- 0:03
δ Hydra et Crateris	+ 0:03
ν Leonis	- 0:04
β Leonis	+ 0:01
β Chamaeleontis	+ 0:38
β Hydry	- 0:10
γ Virginis	- 0:06

JOINT REPORT BY MESSRS. SMALLLEY AND TODD.

We, the undersigned, having been instructed by our respective Governments of New South Wales and South Australia to determine the common boundary line of the said colonies, as defined by Imperial legislation to be the 141st meridian of east longitude, do hereby declare that we have jointly fixed the same by astronomical observations; and we also declare that, starting from the north bank of the river Murray, the said meridian line is about two miles minutes chains east of the prolongation of the present boundary line between Victoria and South Australia (the north end of which is marked by a pile of stones twelve feet high on the south bank of the river Murray), and that its position is permanently indicated by a substantial brick pyramid, built on the scarp forming the limit of the Murray floods, measuring five feet six inches square at the base, and thirteen feet six inches high, having the words "Province Boundary" on the north and south faces, "N.S.W., G. R. Smallley," on the east face, and "S.A., Charles Todd," on the west face; the said mark being situated about seventy yards from the nearest point of the present line of telegraph, and north of the same; the bend of the river Murray immediately to the east of Slaney's Island being nearly three and a half miles distant, on an astronomical bearing of about 53° west of south, Mount Hancock bearing about 16½° east of south.

And we hereby agree, on behalf of our respective Governments, to accept the line hereinbefore described as the common boundary line of the two colonies.


GEORGE R. SMALLLEY,

Government Astronomer for New South Wales.

CHARLES TODD,

Observer and Superintendent of Telegraphs, South Australia.

Adelaide, South Australia, 8th December 1868.



MAP showing the 1850 BOUNDARY

and part of

BOUNDARY

between

South Australia

and

New South Wales

as fixed by

Observations

Charles Todd,
Surveyor & Inst. of Telegraphs

Arthur Cooper
Deputy Surveyor

1 MILE
1 KILOMETER

Scale
1 MILE
1 KILOMETER